

January 01, 2018	The Quantitative Anatomy of the Dorsal Scapholunate Interosseous Ligament	Background: The anatomy of the scapholunate interosseous ligament (SLIL) has been described qualitatively in great detail, with recognition of the dor
September 01, 2019	The efficacy of Ultra-High Frequency Ultrasonography in the diagnosis of intraoral lesions	Objectives: The aim of the present study was to evaluate the diagnostic efficacy of ultra-high frequency ultrasound (UHFUS) imaging of intraoral soft
May 01, 2019	Radial artery remodeling following transradial percutaneous coronary intervention in men and women: insights from serial ultrahigh frequency ultrasonography	Background: Remodeling of the radial artery (RA) after transradial percutaneous coronary intervention (TRI) is under studied.
March 05, 2019	Mechanical Deformation of Human Optic Nerve Head and Peripapillary Tissue in Response to Acute IOP Elevation	PURPOSE.
March 01, 2019	Feasibility of a combination of intraoral UHFUS and CBCT in the study of peri-implantitis	Objectives: The aim of this study was to investigate the combination of intraoral ultra-high-frequency ultrasonography (UHFUS) and cone beam computed

January 01, 2019	Advanced evaluation of hidradenitis suppurativa with ultra-high frequency ultrasound: A promising tool for the diagnosis and monitoring of disease progression	Background: Hidradenitis suppurativa is a chronic inflammatory skin disease.
January 01, 2019	Intraoral Ultra-High Frequency Ultrasound study of oral lichen planus: A pictorial review	Background: Ultra-High Frequency Ultrasound (UHFUS) is a recently introduced diagnostic technique involving the use of higher frequencies compared to
January 01, 2019	Ultra High-frequency Ultrasonographic Imaging with 70 MHz Scanner for Visualization of the Lymphatic Vessels	Background: Identification and localization of functional lymphatic vessels are important for lymphaticovenular anastomosis.
January 01, 2019	Ultra-high-frequency Ultrasound to Assess Nerve Fascicles in Median Nerve Traumatic Neuroma	A traumatic neuroma is a major cause of persistent neuropathic pain. Diagnostic imaging tools are critical to the success of surgical treatment.
January 01, 2019	Comparison of high-frequency and ultrahigh-frequency probes in chronic inflammatory demyelinating polyneuropathy	Objectives: High-frequency ultrasound (HFUS 18–20 MHz) performed on patients with chronic inflammatory demyelinating polyneuropathy (CIDP) shows a foc
January 01, 2019	Advanced ultrasound techniques for pediatric imaging	Ultrasound has become a useful tool in the workup of pediatric patients abstract because of the highly convenient, cost-effective, and safe nature of

<p>December 23, 2018</p>	<p>The 'ALSPAC in London' dataset: adiposity, cardiometabolic risk profiles, and the emerging arterial phenotype in young adulthood</p>	<p>Rising rates of adiposity in the young pose one of the greatest threats to future population burden of cardiovascular disease.</p>
<p>December 20, 2018</p>	<p>Ultrasound Characteristics of the Hair Follicles and Tracts, Sebaceous Glands, Montgomery Glands, Apocrine Glands, and Arrector Pili Muscles</p>	<p>Objectives—To explore the capability of very high-frequency ultrasound (US; 50–71 MHz) to detect the normal morphologic characteristics of the hair fo</p>
<p>December 06, 2018</p>	<p>Preliminary experience of the use of high-resolution skin ultrasound for the evaluation of extrathyroideal manifestations of Graves' disease and response to UVA-1 phototherapy</p>	<p>Graves' orbitopathy (GO) and pre-tibial mixedema (PTM) are autoimmune manifestation sharing the same etiology and histopathology, a chronic course and</p>
<p>November 01, 2018</p>	<p>Motor cortex neurovascular coupling: inputs from ultra-high-frequency ultrasound imaging in humans</p>	<p>OBJECTIVE Neurovascular coupling reflects the link between neural activity and changes in cerebral blood flow.</p>
<p>July 01, 2018</p>	<p>In vivo estimation of the Young's modulus in normal human dermis</p>	<p>Skin elastic properties change during a cutaneous disorder or in the aging process.</p>

<p>June 01, 2018</p>	<p>Ultra-high frequency ultrasound in planning capillary perforator flaps: Preliminary experience☆</p>	
<p>January 01, 2018</p>	<p>High-Frequency Micro-Ultrasound Imaging and Optical Topographic Imaging for Spinal Surgery: Initial Experiences</p>	<p>High frequency micro-ultrasound (μUS) transducers with central frequencies up to 50 MHz facilitate dynamic visualization of patient anatomy with minim</p>
<p>January 01, 2018</p>	<p>Ultrahigh-resolution ultrasound characterization of access site trauma and intimal hyperplasia following use of a 7F sheathless guide versus 6F sheath/guide combination for transradial artery PCI: Results of the PRAGMATIC trial</p>	<p>There exist limited data on the relative degree of acute injury and late healing of the radial artery after transradial artery (TRA) percutaneous coro</p>
<p>January 01, 2018</p>	<p>Very-High-Resolution Sonography Of Median Nerve : A Comparative Study Vs High-Resolution Sonography In Healthy Subjects</p>	<p>The use of ultrasound in the study of peripheral nerves dates back to the late 80s [1]; the first ultrasound studies of the median nerve (MN) affected</p>

January 01, 2018	Intraoperative imaging of lymphatic vessel using ultra high-frequency ultrasound	
January 01, 2018	New findings in non-invasive imaging of cutaneous endometriosis: Dermoscopy, high-frequency ultrasound and reflectance confocal microscopy	© 2018 John Wiley & Sons A/S. Background: Cutaneous endometriosis (CE) is rare and its dermoscopic features were reported only in 3 patients.
January 01, 2018	Ultrahigh Frequency Ultrasound Imaging of the Hand: A New Diagnostic Tool for Hand Surgery	Background: Ultrasonography is a cost-effective, noninvasive, and expedient imaging modality with numerous clinical applications.
April 09, 2017	Clinical and biological markers of premature aging after autologous SCT in childhood cancer	The aim of this study was to analyze the prevalence of frailty and physical health limitations among long-term survivors of high-risk neuroblastoma (H
March 01, 2017	Development of an injectable pseudo-bone thermo-gel for application in small bone fractures	A pseudo-bone thermo-gel was synthesized and evaluated for its physicochemical, mechanical and rheological properties, with its application to treat s
February 01, 2017	Ultra High-frequency Ultrasound of Fascicles in the Median Nerve at the Wrist	Introduction: An ultra high-frequency (70 MHz) ultrasound device has recently been approved for human use.

January 01, 2016	Comparison of very-high-frequency ultrasound assessment of radial arterial wall layers after first and repeated transradial coronary procedures	BACKGROUND Transradial coronary procedure (TRP) traumatizes the radial artery (RA), especially resulting in changes to arterial wall morphology.
January 01, 2016	Computer-Aided Evaluation of Blood Vessel Geometry From Acoustic Images	A method for computer-aided assessment of blood vessel geometries based on shape-fitting algorithms from metric vision was evaluated.
January 01, 2016	Quantification of granuloma volume and response to treatment in cutaneous sarcoidosis using 3-dimensional high-frequency ultrasound scan	C utaneous involvement occurs in 25% to 30% of patients with sarcoidosis. ¹ Effective treat- ment of the disease has historically been challenging give
January 01, 2016	Ultrahigh Frequency Ultrasound Imaging of the Hand : A New Diagnostic Tool for Hand Surgery	Background: Ultrasonography is a cost-effective, noninvasive, and expedient imaging modality with numerous clinical applications.
September 06, 2016	Neonatal Arterial Morphology Is Related to Body Size in Abnormal Human Fetal GrowthCLINICAL PERSPECTIVE	BACKGROUND Restriction in fetal growth is associated with cardiovascular disease in adulthood.
July 01, 2016	Ultra high-frequency ultrasound: New capabilities for nail anatomy exploration	Recent development of ultra high-resolution ultrasound systems, with frequencies as high as 70 MHz and capabil- ity resolution as fine as 30 μ m, could

February 02, 2016	The Rotterdam Radial Access Research	Background—Radial artery wall might be damaged after cannulation for cardiac catheterization.
January 01, 2015	Photoacoustic imaging of real-time oxygen changes in chronic leg ulcers after topical application of a haemoglobin spray: a pilot study	Objective: To use a non-invasive measurement of oxygen saturation in chronic leg ulcers after the application of a topical haemoglobin spray to invest
November 01, 2015	Radiotherapy-related arterial intima thickening and plaque formation in childhood cancer survivors detected with very-high resolution ultrasound during young adulthood	PURPOSE: To test intensive alkylator-based therapy in desmoplastic small round-cell tumor (DSRCT).
January 01, 2014	Radial artery intima-media thickness predicts major cardiovascular events in patients with suspected coronary artery disease	AIMS: In the present study, we investigated the prognostic value of radial artery intima-media thickness (rIMT) in patients with suspected coronary ar
June 01, 2013	High-frequency micro-ultrasound for vascular access in young children--a feasibility study by the High-frequency UltraSound in Kids studY (HUSKY) group.	BACKGROUND: Cannulation of small arteries and veins in young children can be challenging.

<p>February 22, 2013</p>	<p>Increased Rate of Arterial Stiffening with Obesity in Adolescents: A Five-Year Follow-Up Study</p>	<p>BACKGROUND: We prospectively and longitudinally determined the effects of childhood obesity on arterial stiffening and vascular wall changes.</p>
<p>January 14, 2013</p>	<p>Assessment of vascular remodeling after the Fontan procedure using a novel very high resolution ultrasound method: arterial wall thinning and venous thickening in late follow-up</p>	<p>The Fontan circulation is associated with an increased central venous pressure, decreased ventricular preload, and increased afterload.</p>
<p>January 01, 2013</p>	<p>The potential influence of diabetic history on peripheral blood flow in superficial skin</p>	<p>Vascular complication occurrence increases with the duration of diabetes.</p>
<p>March 01, 2012</p>	<p>High-resolution radial artery intima-media thickness and cardiovascular risk factors in patients with suspected coronary artery disease – Comparison with common carotid artery intima-media thickness</p>	<p>Objective: The radial artery wall structure can be measured with non-invasive very high-resolution ultrasound with great feasibility and high accuracy</p>

<p>January 01, 2011</p>	<p>Assessment of early radial injury after transradial coronary intervention by high-resolution ultrasound biomicroscopy: Innovative technology application</p>	<p>BACKGROUND: Transradial coronary intervention (TRI) introduces injury to the radial artery (RA) which will affect repeat transradial coronary procedur</p>
<p>January 01, 2011</p>	<p>Graft vasculopathy in clinical hand transplantation</p>	<p>Allogeneic hand transplantation is now a clinical reality.</p>
<p>December 01, 2011</p>	<p>Feasibility of very-high resolution ultrasound to assess elastic and muscular arterial wall morphology in adolescents attending an outpatient clinic for obesity and lipid abnormalities</p>	<p>Objective: Atherosclerosis begins during early life and is accelerated in individuals with cardiovascular risk factors. We hypothesized that very-high resolution ultr</p>
<p>November 01, 2011</p>	<p>Arteriovenous Fistulas for Hemodialysis: Application of High-Frequency US to Assess Vein Wall Morphology for Cannulation Readiness</p>	<p>To determine whether venous wall thickness and hoop (circumferential) stress, as determined with high-frequency ultrasonography (US), can predict cann</p>
<p>January 01, 2010</p>	<p>Epidermal Thickness and Biomechanical Properties of Plantar Tissues in Diabetic Foot</p>	<p>Diabetic foot is a common complication for people with diabetes but it is unclear whether the change is initiated from the skin surface or underneath</p>

<p>October 01, 2010</p>	<p>Transcutaneous very-high-resolution ultrasound to quantify arterial wall layers of muscular and elastic arteries: Validation of a method</p>	<p>Background: High-resolution ultrasound (HRU) is used to measure carotid intima-media thickness (IMT).</p>
<p>July 01, 2010</p>	<p>High-resolution ultrasound showing increased intima and media thickness of the radial artery in patients with end-stage renal disease</p>	<p>Objective: Although clinically relevant atherosclerosis of the upper limb arteries is rarely seen, intimal hyperplasia of the arteries may reflect glo</p>
<p>January 11, 2010</p>	<p>Thicker carotid intima layer, thinner media layer and higher intima/media ratio in women with recurrent depressive disorders: A pilot study using non-invasive high frequency ultrasound</p>	<p>Objectives: Growing evidence indicates that depression is an important risk factor for coronary heart disease.</p>
<p>January 01, 2009</p>	<p>Increased intima thickness of the radial artery in individuals with prehypertension and hypertension</p>	<p>Background: We have used a novel ultra high-frequency (55 MHz) ultrasound technique to non-invasively measure the radial arterial vessel wall and sepa</p>
<p>December 01, 2009</p>	<p>High-resolution ultrasonography of the cutaneous nerve branches in the hand and wrist</p>	<p>Ultrasonography can be used in the diagnosis of various neuropathies, including nerve injury.</p>

<p>January 01, 2009</p>	<p>High-Frequency Ultrasound for Evaluation of Intimal Thickness</p>	<p>Background: The measurement of carotid intima–medial thickness is a well-validated measure of cardiovascular risk.</p>
<p>May 01, 2008</p>	<p>Limb Stress-Rest Perfusion Imaging With Contrast Ultrasound for the Assessment of Peripheral Arterial Disease Severity</p>	<p>OBJECTIVES: We hypothesized that stress-rest perfusion imaging of skeletal muscle in the lower extremity with contrast-enhanced ultrasound (CEU) could</p>
<p>January 01, 2008</p>	<p>Obese children show increased intimal wall thickness and decreased pulse wave velocity</p>	<p>OBJECTIVE: Childhood obesity confers an increased risk of vascular changes and adult cardiovascular disease.</p>
<p>January 01, 2007</p>	<p>Increasing peripheral artery intima thickness from childhood to seniority</p>	<p>BACKGROUND: Using new, very high-resolution ultrasound biomicroscopy, we examined the thickness of artificial layers of silicone and intima thickness</p>